

# **In-service training for health professionals to improve care of the seriously ill newborn or child in low and middle-income countries (Review) (Review)**

Opiyo N, English M



**THE COCHRANE  
COLLABORATION®**

This is a reprint of a Cochrane review, prepared and maintained by The Cochrane Collaboration and published in *The Cochrane Library* 2010, Issue 5

<http://www.thecochranelibrary.com>



---

In-service training for health professionals to improve care of the seriously ill newborn or child in low and middle-income countries  
(Review) (Review)  
Copyright © 2010 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

## TABLE OF CONTENTS

HEADER . . . . .	1
ABSTRACT . . . . .	1
PLAIN LANGUAGE SUMMARY . . . . .	2
SUMMARY OF FINDINGS FOR THE MAIN COMPARISON . . . . .	2
BACKGROUND . . . . .	5
OBJECTIVES . . . . .	7
METHODS . . . . .	7
RESULTS . . . . .	9
Figure 1. . . . .	10
Figure 2. . . . .	11
Figure 3. . . . .	11
DISCUSSION . . . . .	11
AUTHORS' CONCLUSIONS . . . . .	12
ACKNOWLEDGEMENTS . . . . .	13
REFERENCES . . . . .	13
CHARACTERISTICS OF STUDIES . . . . .	15
DATA AND ANALYSES . . . . .	18
Analysis 1.1. Comparison 1 Opiyo 2008, Outcome 1 Mortality. . . . .	18
Analysis 2.1. Comparison 2 Senarath 2007, Outcome 1 Practice of preparedness of resuscitation. . . . .	19
Analysis 2.2. Comparison 2 Senarath 2007, Outcome 2 Preparedness for resuscitation - adjusted for clustering. . . . .	19
APPENDICES . . . . .	19
WHAT'S NEW . . . . .	32
HISTORY . . . . .	32
CONTRIBUTIONS OF AUTHORS . . . . .	33
DECLARATIONS OF INTEREST . . . . .	33
SOURCES OF SUPPORT . . . . .	33
INDEX TERMS . . . . .	33

[Intervention Review]

# In-service training for health professionals to improve care of the seriously ill newborn or child in low and middle-income countries (Review)

Newton Opiyo<sup>1</sup>, Mike English<sup>1</sup>

<sup>1</sup>Child and Newborn Health Group, Kenya Medical Research Institute/Wellcome Trust Research Programme, Nairobi, Kenya

Contact address: Newton Opiyo, Child and Newborn Health Group, Kenya Medical Research Institute/Wellcome Trust Research Programme, PO Box 43640, Nairobi, 00100 GPO, Kenya. [nopiyo@nairobi.kemri-wellcome.org](mailto:nopiyo@nairobi.kemri-wellcome.org).

**Editorial group:** Cochrane Effective Practice and Organisation of Care Group.

**Publication status and date:** Edited (no change to conclusions), published in Issue 5, 2010.

**Review content assessed as up-to-date:** 21 May 2009.

**Citation:** Opiyo N, English M. In-service training for health professionals to improve care of the seriously ill newborn or child in low and middle-income countries (Review). *Cochrane Database of Systematic Reviews* 2010, Issue 4. Art. No.: CD007071. DOI: 10.1002/14651858.CD007071.pub2.

Copyright © 2010 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

## ABSTRACT

### Background

A variety of emergency care training courses based on developed country models are being promoted as a strategy to improve the quality of care of the seriously ill newborn or child in developing countries. Clear evidence of their effectiveness is lacking.

### Objectives

To investigate the effectiveness of in-service training of health professionals on their management and care of the seriously ill newborn or child in low and middle-income settings.

### Search strategy

We searched The Cochrane Register of Controlled Trials (CENTRAL), the Specialised Register of the Cochrane EPOC group (both up to May 2009), MEDLINE (1950 to May 2009), EMBASE (1980 to May 2009), CINAHL (1982 to March 2008), ERIC / LILACS / WHOLIS (all up to October 2008), and ISI Science Citation Index Expanded and ISI Social Sciences Citation Index (both from 1975 to March 2009). We checked references of retrieved articles and reviews and contacted authors to identify additional studies.

### Selection criteria

Randomised controlled trials (RCTs), cluster-randomised trials (CRTs), controlled clinical trials (CCTs), controlled before-after studies (CBAs) and interrupted time series studies (ITSs) that reported objectively measured professional practice, patient outcomes, health resource /services utilization, or training costs in healthcare settings (not restricted to studies in low-income settings).

### Data collection and analysis

We independently selected studies for inclusion, abstracted data using a standardised form, and assessed study quality. Meta-analysis was not appropriate. Study results were summarised and appraised.

---

**In-service training for health professionals to improve care of the seriously ill newborn or child in low and middle-income countries (Review) (Review)**

Copyright © 2010 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

I

## **Main results**

Two studies of varied designs were included. In one RCT of moderate quality, Newborn Resuscitation Training (NRT) was associated with a significant improvement in performance of adequate initial resuscitation steps (risk ratio 2.45, 95% confidence interval (CI) 1.75 to 3.42,  $P < 0.001$ , adjusted for clustering) and a reduction in the frequency of inappropriate and potentially harmful practices (mean difference 0.40, 95% CI 0.13 to 0.66,  $P = 0.004$ ). In the second RCT, available limited data suggested that there was improvement in assessment of breathing and newborn care practices in the delivery room following implementation of Essential Newborn Care (ENC) training.

## **Authors' conclusions**

There is limited evidence that in-service neonatal emergency care courses improve health-workers' practices when caring for a seriously ill newborn although there is some evidence of benefit. Rigorous trials evaluating the impact of refresher emergency care training on long-term professional practices are needed. To optimise appropriate policy decisions, studies should aim to collect data on resource use and costs of training implementation.

## **PLAIN LANGUAGE SUMMARY**

### **Effectiveness of in-service training in the care of the seriously ill newborn or child**

In developing countries, most deaths in very ill babies and children who seek care in healthcare facilities happen within 48 hours of being seen. Currently, a number of emergency care courses, adapted from developed countries are being promoted in developing countries as a means to improving the quality of care provided to seriously ill newborns or children. Whether these courses result in improvement in health workers' ability to provide appropriate care remains unclear.

Although the results from the two included studies showed that emergency care training could be followed by improvement in health workers' practices, because of the small number of studies, differences in the training courses, and weaknesses in the study methods, it is not possible to conclude that in general such in-service training improves health worker practices when they are faced with a seriously ill child. Further well-conducted studies are therefore needed to provide reliable evidence on what such courses achieve. To guide decisions regarding which interventions to invest in, such studies should also collect data on resources used and costs of training interventions.

## SUMMARY OF FINDINGS FOR THE MAIN COMPARISON [\[Explanation\]](#)

In-service neonatal emergency care training versus standard care for healthcare professionals						
<b>Patient or population:</b> Healthcare professionals <b>Settings:</b> Hospital-based settings <b>Intervention:</b> In-service neonatal emergency care training <b>Comparison:</b> Usual or standard care						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Usual or standard care	In-service neonatal emergency care training				
<b>Proportion of appropriate initial resuscitation steps</b> Direct observations Follow-up: 50 days			RR 2.45 (1.75 to 3.42)	83 [1 study]	++ +0 moderate	Overall quality: high risk of bias
	27 per 1000	67 per 100 (48 to 93)				
	Medium risk population					
<b>Frequency of inappropriate practices</b> Follow-up: mean 50 days	The mean frequency of inappropriate practices in the control group was 0.92	The mean frequency of inappropriate practices in the control group was <b>0.40 higher</b> (0.13 to 0.66 higher)		83 [1 study]	++ +0 moderate	
<b>Mortality</b> Direct observations Follow-up: mean 40 days			RR 0.77 (0.40 to 1.48)	90 (1 study)	++ +0 moderate	Trial not sufficiently powered to detect a mortality effect
	36 per 100	28 per 100 (14 to 53)				
	Medium risk population					

\*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; RR: Risk Ratio

GRADE Working Group grades of evidence

**High quality:** Further research is very unlikely to change our confidence in the estimate of effect.

**Moderate quality:** Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

**Low quality:** Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

**Very low quality:** We are very uncertain about the estimate.

## BACKGROUND

In developing countries most deaths among seriously ill children who come into contact with referral level health services occur within 48 hours of being seen (Berkley 2005). It is possible that good quality immediate and effective care provided by health professionals could reduce these deaths (Nolan 2001). Provision of appropriate care is, however, dependent on the presence of adequately skilled health personnel at the point of delivery (WHO 2005). To improve health workers' capacity to provide effective care for seriously ill children and newborns in low-income countries, a number of in-service training courses, mainly based on developed countries' models, are proposed.

These courses include: (1) neonatal life support courses (e.g. Newborn Life Support (NLS), Neonatal Resuscitation Program (NRP)), (2) paediatric life support courses (e.g. Paediatric Advanced Life Support (PALS), Paediatric Life Support (PLS)), (3) life support / emergency care elements within the Integrated Management of Pregnancy and Childbirth (e.g. Essential Newborn Care (ENC)) and (4) components of other in-service child health training courses that deal with care of serious illness (e.g. Emergency Triage, Assessment and Treatment (ETAT), Control of Diarrheal Diseases (CDD) and Acute Respiratory Infections (ARI)

case management programs and the training components of the Integrated Management of Childhood Illness (IMCI) strategy).

Although such formalised educational programs vary in origin, scope and target audience, they are typically aimed at in-service rather than pre-service training, and are short and intensive with a structured approach to the presentation of their clinical subject (Table 1). The one-day NRP course was first taught in 1987 in the USA while the one-day NLS course was initiated in the UK in 2001 (Raupp 2007). PALS, a two-day course, was piloted in USA in 1988. Advanced Paediatric Life Support (APLS), a three-day course, was developed and piloted in the UK in 1992. Two other courses - the one-day PLS course and Prehospital PLS - have been designed to complement the APLS (Jewkes 2003). The World Health Organization (WHO) has recently added to this list the 3½-day ETAT course based upon and validated against the APLS course in Malawi (Gove 1999; Molyneux 2006). This course is aimed specifically at lower income settings and is intended to improve prompt identification and institution of life saving emergency treatment for very ill children. These life support courses emphasize early recognition of neonatal/paediatric emergencies and prevention of cardio-respiratory arrest (and mortality) through resuscitation.

**Table 1. Summary of neonatal and paediatric emergency care courses†**

Course	Subject	Duration (days)	Target audience
Neonatal Life Support (NLS)	Neonatal resuscitation	1	Midwives Paediatricians General Practitioners
Neonatal Resuscitation Program (NRP)	Neonatal resuscitation	1	Midwives Paediatricians General practitioners
Paediatric Life Support (PLS)	Basic Life Support (BLS) and Advanced Life Support (ALS) for children Recognition of paediatric emergencies	1	Nurses and doctors involved in paediatric care
Paediatric Advanced Life Support (PALS)	BLS and ALS for children Recognition of paediatric emergencies Some neonatal life support	2	Nurses and doctors involved in paediatric care
Prehospital Paediatric Life Support (PHPLS)	Prehospital paediatric emergency care	2+	General practitioners, paramedics, some nurses, emergency medicine staff

**Table 1. Summary of neonatal and paediatric emergency care courses†** (Continued)

Advanced Paediatric Life Support (APLS)	BLS and ALS for children Paediatric emergencies, including serious illness and major trauma, some neonatal life support	3	Paediatricians, emergency medicine doctors, some anaesthetists, senior paediatric nurses
Emergency Triage Assessment and Treatment (ETAT)	Very ill children presenting to hospital	3.5	Doctors, nurses, paramedics
Essential Newborn Care (ENC) course	Aspects of newborn care (including neonatal resuscitation) in the Integrated Management of Pregnancy and Childbirth (IMPAC)	5	Nurses, midwives, doctors
Integrated Management of Childhood Illness (IMCI)	Ill children and neonates including emergency care or identification and referral of the seriously ill	11	Nurses, midwives, doctors

†Tulloch 1999, Jewkes 2003, Mello 2003, Irimu 2008

The more general CDD and ARI programs were developed by the WHO in 1980, in recognition of the high childhood mortality due to diarrhea/dehydration and pneumonia for the very ill neonate or child and focus on case management training rather than life-support (Forsberg 2007; Pio 2003). While these courses concentrate predominantly on community or outpatient based management, where there is good evidence for their success (Sazawal 2001), they also include guidance on management of very severe illness. These disease-specific training approaches were incorporated into the broader package of the IMCI strategy. Here the particular focus for management of the very ill child is the decision to refer to hospital and provide pre-referral management. In addition to this, the WHO has developed a specific five-day course on hospital management of severe malnutrition (WHO 2002).

In-service training, however, costs both time and money: for example, the cost of the 2-day European Paediatric Life Support (EPLS) course is estimated to be about USD 190 per trainee in Kenya (Personal communication with ME, 2009). These costs include allowances for the trainers (e.g. travel refunds), course materials (e.g. course manuals, consumable teaching aides, etc) and hotel costs for the participants. Apart from the sometimes high costs of providing courses (recovered in high income countries often with high course fees), attendance at such courses often means that important staff (instructors and participants) are absent from their normal duties with potential disruption to patient care and for

some a loss of personal income (Jabbour 1996). Despite their cost, however, emergency care courses are a thriving enterprise in many high income countries, as reflected in their ever increasing number and variety (Jewkes 2003). In the hope that they might improve the quality of care in many low- and middle-income countries, considerable global efforts and investments have gone into their further development, refinement and adaptation to meet individual country needs (Baskett 2005). Yet despite these investments and the faith placed in them by many organizations and institutions, clear evidence of the effectiveness of these courses in improving health workers ability to manage seriously ill children or neonates appears lacking.

Two Cochrane reviews from the Injuries Group have examined the effectiveness of Advanced Trauma Life Support course (ATLS), for ambulatory crews (Sethi 2001) and hospitals (Shakiba 2003), respectively, in reducing mortality and morbidity for victims of trauma of any age. The pre-hospital review (Sethi 2001) identified one small randomised controlled trial (n = 16) while the hospital review (Shakiba 2003) did not find any relevant randomised controlled trials. The Cochrane review authors concluded that there is no clear evidence that ATLS training impacts on the outcome of victims of trauma.

The effectiveness of in-service training of health professionals depends on changes in health worker practices which, plausibly,



should precede any impact on mortality or morbidity. This review investigated if there are systematic differences in health workers' professional practices (i.e. more appropriate management or referral of seriously ill children/newborns or both), patient outcomes (mortality and morbidity), or health resource use (e.g. drug use, laboratory tests) and services utilization (e.g. length of hospitalization, return visits) after in-service training in emergency care or care for the seriously ill newborn or child. Information regarding the effectiveness of such in-service training courses is required to enable low-income countries to prioritise the health interventions they invest in.

## OBJECTIVES

To investigate the effectiveness of in-service training of health professionals on their management and care of seriously ill neonates or children in low-income settings.

We considered the comparisons listed below.

1. Interventions in which seriously ill neonates have been cared for by qualified health professionals who have undergone neonatal emergency care training compared to those receiving usual or standard care.
2. Interventions in which seriously ill children have been cared for by qualified health professionals who have undergone paediatric emergency care training compared to those receiving usual or standard care.
3. Interventions in which seriously ill neonates or children have been cared for by qualified health professionals who have undergone any other in-service child health training that deals with care of severe illness (e.g. CDD, ARI, ETAT, etc) compared to those receiving usual or standard care.

## METHODS

### Criteria for considering studies for this review

#### Types of studies

All randomised controlled trials (RCTs), cluster-randomised trials (CRTs), controlled clinical trials (CCTs), controlled before-after studies (CBAs) and interrupted time series studies (ITSs) that have evaluated the effects of in-service training on at least one of the outcomes listed below. We did not consider before and after studies that had no parallel control groups.

#### Types of participants

All qualified healthcare professionals, including, doctors (general practitioners and specialists), nurses, pharmacists and dietitians/nutritionists, in outpatient or hospital-based settings, responsible for the management and care of seriously ill neonates or children. We excluded non-qualified healthcare providers (e.g. medical students/trainees, medical interns, community health workers). Studies were not excluded based on the setting (low or high income).

#### Types of interventions

We considered implementation studies of the following in-service training courses aimed at changing provider behaviour in the care of the seriously ill newborn or child:

- 1) Neonatal life support courses e.g. NLS, NRP, and others.
- 2) Paediatric life support courses e.g. PALS, PLS, and others.
- 3) Life support elements within the Integrated Management of Pregnancy and Childbirth e.g. ENC.
- 4) Other in-service newborn and child health training courses aimed at the recognition and management of the seriously ill child e.g. ETAT, CDD, ARI, malaria case management or the training components of the IMCI strategy.

We excluded studies of complex interventions in which training is combined with and impossible to separate from additional health system improvements (for example improved drug/equipment/staff supply/health facility reorganisation).

#### Types of outcome measures

##### Primary outcomes

We included studies only if they reported at least one of the following objectively measured professional (in practice) performance outcomes.

- Adherence to treatment guidelines
- Prescribing practices
- Clinical assessment and diagnosis
- Recognition of and management or referral of the seriously ill newborn/child

##### Secondary outcomes

Where reported, we also considered the following outcomes

- Health resource utilization, for example, use of drugs, laboratory tests, etc.
- Health services utilization, for example, length of hospital stay.
- 'Other markers' of clinical performance, for example, simulated health worker performance (in practice)
- Cost of training, for example, costs of purchasing training materials.

We excluded studies that only reported 'other markers' of performance (for example, simulations/skill testing that is done outside of the practice setting (in the classroom) that are tests of skill, such as practicing/demonstrating resuscitation techniques using a dummy). However, we considered for inclusion simulations of emergency care in the practice setting that were designed to reflect real practice.

## Search methods for identification of studies

See: Cochrane Effective Practice and Organisation of Care Group methods used in reviews.

We selected studies according to the Cochrane Handbook for Systematic Reviews of Interventions (Higgins 2006) and the Effective Practice and Organisation of Care Group (EPOC) methods used in reviews (EPOC 2007).

## Electronic searches

To identify potential studies for inclusion, we searched the following electronic databases.

- a) The Cochrane Central Register of Controlled Trials (CENTRAL)/EPOC register (up to May 2009) (Appendix 1).
- b) MEDLINE (1950 to May 2009), EMBASE (1980 to May 2009), CINAHL (1982 to March 2008), LILACS (up to October 2008), ERIC (up to October 2008) and WHOLIS (up to October 2008). A forward search for papers that cite included studies was conducted in the ISI Science Citation Index Expanded (SCI- Expanded) (1975 to March 2009) and ISI Social Sciences Citation Index (SSCI) (1975 to March 2009).

We developed search strategies for electronic databases using the methodological component of the EPOC search strategy combined with selected MeSH terms and free text terms. Appendix 2 shows the terms used in the MEDLINE search strategy. We modified this search strategy as appropriate for other databases (Appendix 3; Appendix 4; Appendix 5; Appendix 6; Appendix 7). No date or language restrictions were applied.

## Searching other resources

- a) List of references from the Health Care Provider Performance (HCPP) Review (Rowe 2008, available from Alexander K. Rowe, e-mail: axr9@cdc.gov).
- b) Clinical trial registries for ongoing studies
- c) Reviewed reference lists of all papers and relevant reviews identified.
- d) Contacted authors of relevant articles regarding any further published or unpublished work.

## Data collection and analysis

### Selection of studies

The two review authors independently screened the titles and abstracts (where available) based on the pre-determined review criteria. We retrieved all full text copies of studies meeting the inclusion criteria for a detailed assessment by both authors. Disagreements were resolved through consensus following discussion between the authors.

### Data extraction and management

Both authors independently extracted data from trial reports using a modified EPOC data collection tool from the EPOC group (EPOC 2007). We extracted data relating to the following items:

1. Participants (healthcare providers and patients): profession, number, age, years of experience, and number of episodes/practices performed by the included healthcare providers. We also collected data regarding the number and specific clinical problems of the included patients.
2. Intervention: type and duration of training and co-interventions (teachings aids, self-learning manuals, etc).
3. Type of targeted behaviour (general management of the problem).
4. Study designs and the key characteristics of the studies (setting, unit of allocation/analysis, length of post-intervention follow-up, and time lag between the intervention and post-intervention assessment).
5. Results grouped according to the primary and secondary outcomes specified above.

We resolved disagreements through discussion.

### Assessment of risk of bias in included studies

Both review authors independently assessed the risk of bias of all included studies using the EPOC checklist for the assessment of methodological quality of studies (see EPOC module) and rated them into three classes: low (low risk of bias for all key domains), high (high risk of bias for one or more key domains) and unclear risk of bias (unclear risk of bias for one or more key domains) based on the assessment of the following: allocation sequence generation, allocation concealment, measurement of baseline outcomes, baseline characteristics of providers, blinding (participants, personnel and outcome assessors), completeness of follow-up (mainly related to follow-up of professionals), treatment of incomplete outcome data, and protection against selective outcome reporting and contamination. We resolved disagreements regarding the quality ratings through discussion between the two authors. Studies were not excluded based on their risk of bias. We assessed the overall quality of evidence using the Grading of Recommendations Assessment,

Development and Evaluation (GRADE) approach, see [Summary of findings for the main comparison](#).

## Data synthesis

Identified eligible studies varied in design, focus (newborn resuscitation, essential newborn care), length (1 versus 4 days) and outcome measured. It was therefore not appropriate to combine the results of the studies. The results are therefore presented separately.

In one study ([Senarath 2007](#)), data analyses could have been improved: First, there was a 'unit of analysis error' - the unit of randomisation was the hospital but the unit of analysis was observed practices at delivery. Thus, the strength of the reported associations could have been over-estimated ('spuriously low P values') as correlation within hospitals was not taken into account. Secondly, comparisons were made within (intervention and control) groups before and after the intervention but effects in the experimental and control groups were not directly compared. Re-analysis was, however, only possible for the outcome on preparedness for resuscitation - where baseline levels of performance ('mean practice scores') were comparable (intervention group  $7.0 \pm 4.08$  versus control group  $7.21 \pm 4.51$ ). The re-analysis involved comparison of intervention and control groups through computation of a mean difference using the reported standard deviations to estimate standard errors. To account for clustering we assumed an intra-cluster correlation coefficient (ICC) of 0.015 (with a design effect of 1.129), based on published data ([Rowe 2002](#)). The re-calculated P value was annotated with 're-analysed'.

## RESULTS

### Description of studies

See: [Characteristics of included studies](#); [Characteristics of excluded studies](#).

See: [Characteristics of included studies](#); [Characteristics of excluded studies](#).

In total, we identified 2480 references from both the electronic and supplementary searches. No ongoing studies were identified. Not all the identified articles were published in English: 137 titles/abstracts (from the LILAC database) were in Portuguese and their titles were translated to English. We excluded 2334 irrelevant articles following a review of all the titles and abstracts. Reasons for exclusion included: inadequate study designs, inappropriate interventions/outcomes, enrolment of non-qualified healthcare providers, assessment of simulated practices outside practice settings, letters to the editor, commentaries, review articles, guidelines, non-paediatric studies, etc. We retrieved the full texts of 146 papers for further eligibility assessment. From these, we identified

eight studies as potentially meeting the review inclusion criteria. We excluded six of these studies for a variety of reasons following a detailed assessment (see [Characteristics of excluded studies](#)): [Bryce 2005](#), a non-randomised controlled study on health facility IMCI training, was excluded as the training intervention was combined and impossible to separate from concurrent district health strengthening activities such as skills reinforcement through supervised clinical practice (i.e. a complex intervention). [El-Arifeen 2004](#), a CRT on the effect of IMCI training on quality of care was excluded as data on referral rate (the appropriate health-worker response to an encounter with a seriously ill child, and our outcome of interest) were not reported for very ill children. We excluded another study ([Gouws 2004](#)) on the effect of IMCI on health worker antibiotic use as no baseline assessment of outcomes was performed in the IMCI trained and untrained groups. One intervention study ([Nadel 2000](#)) of periodic mock resuscitations combined with an 8-hour resuscitation course was excluded as it lacked a concurrent comparison group (i.e. used a historical control group). Two further studies were excluded as they enrolled only apparently well children ([Pelto 2004](#)) or those with mild ARI episodes ([Ochoa 1996](#)). Overall, we have considered two studies that met all the review inclusion criteria (see [Characteristics of included studies](#)). As a formal meta-analysis was not possible - given the small number of studies and differences in interventions (content, format, timing) and reported outcomes - a description of the included studies is provided below.

Both of the included studies were set in the delivery room/theatre in low-income countries (Kenya ([Opiyo 2008](#)), Sri Lanka ([Senarath 2007](#))). Both of the included studies were RCTs. The health professionals were nurses in one trial ([Opiyo 2008](#)) and mixed (doctors, nurses, midwives) in another ([Senarath 2007](#)). The targeted behaviours were process of initiating newborn resuscitation ([Opiyo 2008](#)) and general management/preparation and conduct of delivery care for newborns ([Senarath 2007](#)). The length of time during which the intervention was measured after initiation of intervention was 50 days in [Opiyo 2008](#), and three months in [Senarath 2007](#).

The number of experimental and control groups was balanced in [Opiyo 2008](#) but not in [Senarath 2007](#), where two hospitals were allocated to the intervention group and three hospitals to the control group. Both of the included studies were adequately powered (90%) for the primary outcomes. The unit of allocation in the included studies was healthcare professionals ( $n = 83$ ) ([Opiyo 2008](#)), and hospitals ( $n = 5$ ) ([Senarath 2007](#)). Both dichotomous (for example proportion of adequate resuscitation steps, proportion of newborns with undesirable health events) and continuous outcomes (for example frequency of harmful practices, mean scores of ENC practices) were considered in the included studies. None of the studies included information on the impact of the interventions on healthcare costs or resource utilization.

The first study ([Opiyo 2008](#)) was a RCT to determine if a simple one day newborn resuscitation training (NRT) alters initial

health worker resuscitation practices in a public hospital setting in Kenya. The intervention was a 1-day newborn resuscitation course adapted from the approach of the UK Resuscitation Council. The course teaches an A(Airway), B(Breathing), and C(Circulation) approach to resuscitation laying down a clear step by step strategy for the first minutes of resuscitation at birth. The teaching strategy was comprised of focused lectures and practical scenario sessions using infant manikins. Candidates were provided with a simple instruction manual two weeks before the training for self-learning. Health workers were randomly allocated to receive early training (n = 28) or late training (the control group, n = 55). Data were collected on 97 and 115 resuscitation episodes over 7 weeks after early training in the intervention and control groups respectively. The second study (Senarath 2007) was a RCT (with random allocation to groups) to evaluate the effectiveness of training for care providers on practice of essential newborn care in hospitals in Sri Lanka. The intervention was a 4-day training program on essential newborn care based on the WHO *Training Modules on Essential Newborn Care and Breastfeeding*. Additionally, participants were provided with teaching aids on *Newborn Care* (adapted from the National Neonatology Forum India) and *Resuscitation of the Newborn* (adapted from the Resuscitation Council (UK)). The teaching strategies involved lecture discussions, demonstrations, hands-on training, practical assignments, and small group discussions. Hospitals were randomly assigned to either the intervention group (n = 2 hospitals) or control group (n = 3 hospitals). The main sample for data collection by exit interview included 446 mother-newborn pairs pre-intervention and 446 post-intervention (223 each in intervention and control groups). These exit interview data were not relevant to the topic of this review. Direct observations of delivery practices were however made on a sub-sample consisting of 96 participants (48 before and 48 after the intervention). Post-intervention data collection commenced three months after the intervention.

## Risk of bias in included studies


Both the included studies were of inadequate quality (high risk of bias) (see [Risk of bias in included studies](#)). In Opiyo 2008, blinding of outcome assessors and follow-up of providers was done, while allocation sequence generation, concealment, baseline measurement (of primary outcome), reporting of the reliability of outcome measures, and protection against contamination were not clear. In Senarath 2007, random allocation was adequately concealed, there was complete reporting of outcome data, and the study was adequately protected against contamination and selective outcome reporting. However, allocation sequence generation was unclear, and there were baseline differences in appropriate essential newborn care practices, and in the characteristics of study and control providers. Also, outcomes of interest were not assessed blindly and the presence of a 'unit of analysis error' could have contributed to additional risk of bias.

## Effects of interventions

See: [Summary of findings for the main comparison](#) [Summary of Findings table](#)


In Opiyo 2008, trained providers demonstrated a higher proportion of adequate initial resuscitation steps compared to the control group (trained 66% versus control 27%; risk ratio 2.45, 95% confidence interval (CI) 1.75 to 3.42,  $P < 0.001$ , adjusted for clustering). In addition, there was a statistically significant reduction in the frequency of inappropriate and potentially harmful practices per resuscitation in the trained group (trained 0.53 versus control 0.92, mean difference 0.40, 95% CI 0.13 to 0.66,  $P = 0.004$ ). Group comparison for the overall mortality in all resuscitation episodes (reported but not a stated primary outcome) showed no statistically significant differences between the groups (trained 0.28 (18/65), 95% CI 0.17 to 0.40; control 0.25 (9/25), 0.12 to 0.42,  $P = 0.77$ ) (Figure 1).

**Figure 1. Forest plot of comparison: 2 Opiyo 2008, outcome: 2.1 Mortality.**




In [Senarath 2007](#), assessment of breathing of the newborn at birth and four out the five components of ENC practices improved in the intervention group three months after the intervention. Apart from the outcome on preparedness for resuscitation (see section on data synthesis above), it was not possible to re-analyse the data on other outcomes of interest. In the re-analysis accounted for clustering, ENC course was associated with a significant improvement in resuscitation preparedness (mean difference, MD 8.83, 95% CI 6.41 to 11.25, P value (re-analysed) < 0.001) ([Figure 2](#), [Figure 3](#)).

**Figure 2. Forest plot of comparison: I Senarath 2007, outcome: I.1 Practice of preparedness of resuscitation.**



**Figure 3. Forest plot of comparison: I Senarath 2007, outcome: I.2 Preparedness for resuscitation - adjusted for clustering.**



## DISCUSSION

This review found few well-conducted studies on the impact of neonatal or paediatric in-service training aimed at improving care for the seriously ill newborn or child. Limited evidence from the two included studies suggests a beneficial effect in the following outcomes: performance of initial resuscitation practices and reduction in the frequency of inappropriate practices ([Opiyo 2008](#)) in the short-term, and delivery room newborn care practices ([Senarath 2007](#)). We found no evidence of an effect on mortality, although the only study that reported this outcome was under-

powered to detect a mortality effect. Even though both the included studies reported positive performance outcomes following successful training, a generalisable evidence of effectiveness cannot be inferred - given the differences in interventions, outcomes, clinical settings and weaknesses in the study methods. These results are therefore intended for descriptive purposes only.

The common trend of benefit in the included studies should be interpreted with caution. First, in the study by [Opiyo 2008](#), assessments were conducted immediately following training for a short period of 50 days. Instantaneous improvement in performance would therefore have been expected. Clinical skills have however been shown to decay over time, with as much as a 50%

reduction in performance (as assessed in classroom simulations) within six months of intense training (McKenna 1985). Thus, evaluation of potential interaction effects of training over-time would have produced more reliable results. Conversely, the potential for a 'decay effect' underscores the need for periodic refresher trainings as a means to maintaining optimal performance especially given the infrequent nature of emergency care. Second, in Senarath 2007, baseline performance of newborn care practices was relatively high in both the intervention and control groups. Thus, the narrow 'performance improvement gap' could have limited possible demonstration of a real impact of the ENC program (i.e. possible 'ceiling effect'). Third, training coverage was low in Opiyo 2008 and unclear in Senarath 2007. Saturation training to the level of that reported in one excluded study (El-Arifeen 2004 (94%)) can potentially create a 'herd effect' on provider practices. Thus, possible mediation of reported effects by differences in levels of training coverage cannot be excluded.

The lengths of the considered training interventions varied: 1-day NRT in Opiyo 2008, and 4-day ENC course in Senarath 2007. Apart from the clear effect on costs, there is some evidence that the duration of training courses could influence their effectiveness: one related review (Rowe 2008) (n = 2 studies) which compared the standard IMCI training (duration  $\geq 11$  days) to shortened training (5-11 days) reported marginal effectiveness of the standard in-service IMCI training course over the shortened training. In the same review, the effect of IMCI training over time was mixed with some analyses indicating increased effect with time, while others showed decreasing or no effect. In the current review, the length of follow-up period following training was relatively short (50 days) in Opiyo 2008 and three months in Senarath 2007. Thus, no reliable inference could be made regarding the magnitude of training effect over time. Still, to take account of the potential deterioration of clinical skills over time, it is recommended that evaluations of educational interventions include a sufficient length of follow-up period following the intervention. The effect of training could vary depending on the susceptibility of the targeted behaviour to the training intervention. Some behaviours (such as performance of inappropriate practices e.g. holding the baby upside during resuscitation) are easier to change than others (such as correct performance of all resuscitation steps). In Opiyo 2008, the teaching strategy consisted of focused lectures and practical scenario sessions using an infant manikin, while in Senarath 2007, the strategy involved lecture discussions, demonstrations, hands-on training, practical assignments, and small group discussions. The content and format of in-service training courses could influence their effectiveness - in one Cochrane review on the effects of educational meetings on professional practice and healthcare outcomes (Forsetlund 2009), combined lectures and small group discussions appeared to be more effective. The outlined possible mediators of training effects add to the difficulty in deriving even a qualitative interpretation of the presented evidence.

The limited evidence available can be attributed to a number of factors: First, a significant number of studies were excluded on the basis of inadequate designs (e.g. lack of concurrent controls, use of historical controls, retrospective surveys, naturalistic designs, etc). Thus, the available evidence is mainly of poor quality with unreliable findings. Second, the lack of rigorous trials could also be attributed to design and ethical challenges inherent in the evaluation of educational interventions. Such desirable attributes as protection against contamination cannot practically be achieved within routine practice settings. Random assignment of healthcare providers and already vulnerable populations of infants to a control arm and observation of practices performed by untrained providers clearly raises ethical concerns. Third, effective sample sizes will always be hard to achieve for example severe illness episodes and resuscitation events remain relatively uncommon events in most clinical settings. Thus, large pragmatic multi-centre studies with prolonged observation periods would be needed to sufficiently demonstrate plausible changes in provider performance and ideally mortality. Apart from the clear logistical and cost implications, such trials would have to contend with the difficulty in securing the attendance and continued availability and participation of health workers. A possible optimal design to deal with the above tension between the need for high quality randomised evidence of effectiveness of emergency care courses and the highlighted ethical and practical constraints would be a pragmatic cluster-randomised trial with process evaluations to facilitate a better understanding of the determinants of actual practice (Elie 2007).

The findings of this review, in common with previous related reviews (Jabbar 1996, Rowe 2008), demonstrate the sparse evidence base for the impact of neonatal and paediatric courses on care of the seriously ill newborn or child. None of the included studies considered training programme development or implementation costs and thus any consideration of costs and benefits is impossible. While courses with a broader scope may have a broader range of benefits too there is limited evidence of effectiveness of emergency care courses. However, these courses continue to be popular within ministries of health and healthcare institutions, and are increasingly being promoted by influential groups such as the WHO and its partners. Before these become the standard of care, making them even more difficult to evaluate, evidence of their ability at least to change health worker practices and ideally to reduce mortality are required.

## AUTHORS' CONCLUSIONS

### Implications for practice

Studies included in this review do not provide a definitive evidence of effectiveness of in-service neonatal and paediatric courses in the emergency care setting. Additionally, despite the weak but positive evidence of benefit, it is still uncertain whether such in-service training, compared to alternative interventions, improves



outcomes at reasonable costs. The current findings cannot therefore be used to inform decisions on whether to invest in in-service emergency care training as opposed to other alternatives to improving the survival of seriously ill newborns or children.

## Implications for research

Rigorous trials (with appropriate controls and adequate randomisation procedures) evaluating the impact of refresher emergency care training on long-term outcomes (professional practices and patient outcomes) are needed (given the current uncertainty on how long short-term benefits are retained, particularly in settings where they are used relatively infrequently). Such trials should: 1) involve direct head-to-head comparisons of courses with varied lengths (such as 1-day courses versus 4-day courses); 2) aim to include seriously ill newborns (in out-patient settings) and children (in both out-patient and hospital settings); and 3) include data on resources and cost of training implementation (to optimise appropriate policy decisions regarding which interventions to invest in). To facilitate replication, the studies should also provide sufficient detail regarding their content (e.g. need for equipment, teamwork

and format (e.g. small group interactive versus lectures, hands-on skills with dummies).

## ACKNOWLEDGEMENTS

Newton Opiyo was awarded a Reviews for Africa Programme Fellowship ([www.mrc.ac.za/cochrane/rap.htm](http://www.mrc.ac.za/cochrane/rap.htm)), funded by a grant from The Nuffield Commonwealth Programme, through The Nuffield Foundation.

We would like to thank Marit Johansen and Kjetil Olsen for their assistance with the searches and the acquisition of full text articles. We would also like to thank Andy Oxman and Sasha Shepperd for their inputs at various stages of the review, Alexander Rowe for his helpful comments on the review manuscript, and Susan Babigumira for her editorial assistance.

Mike English is funded by a Wellcome Trust Senior Fellowship (#076827) and Newton Opiyo is supported by funding from a Wellcome Trust Strategic Award (#084538).

## REFERENCES

### References to studies included in this review

#### Opiyo 2008 *{published data only}*

Opiyo N, Were F, Govedi F, Fegan G, Wasunna A, English M. Effect of newborn resuscitation training on health worker practices in Pumwani Hospital, Kenya. *PLoS ONE* 2008;**13**(2):e1599.

#### Senarath 2007 *{published data only}*

Senarath U, Fernando DN, Rodrigo I. Effect of training for care providers on practice of essential newborn care in hospitals in Sri Lanka. *Journal of Obstetric, Gynecologic and Neonatal Nursing* 2007;**36**(6):531–41.

### References to studies excluded from this review

#### Bryce 2005 *{published data only}*

Bryce J, Gouws E, Adam T, Black RE, Schellenberg JA, Manzi F, et al. Improving quality and efficiency of facility-based child health care through Integrated Management of Childhood Illness in Tanzania. *Health Policy and Planning* 2005;**20**(Suppl 1):i69–i76.

#### El-Arifeen 2004 *{published data only}*

El Arifeen S, Blum LS, Hoque DM, Chowdhury EK, Khan R, Black RE, et al. Integrated Management of Childhood Illness (IMCI) in Bangladesh: early findings from a cluster-randomised study. *Lancet* 2004;**364**(9445):1595–602.

#### Gouws 2004 *{published data only}*

Gouws E, Bryce J, Habicht JP, Amaral J, Pariyo G, Schellenberg JA, et al. Improving antimicrobial use among health workers in first-level facilities: results from the multi-country evaluation of the Integrated Management of Childhood Illness strategy. *Bulletin of the World Health Organization* 2004;**82**(7):509–15.

#### Nadel 2000 *{published data only}*

Nadel FM, Lavelle JM, Fein JA, Giardino AP, Decker JM, Durbin DR. Teaching resuscitation to pediatric residents: the effects of an intervention. *Archives of Pediatrics and Adolescent Medicine* 2000;**154**(10):1049–54.

#### Ochoa 1996 *{published data only}*

González Ochoa E, Armas Pérez L, Bravo González JR, Cabrales Escobar J, Rosales Corrales R, Abreu Suárez G. Prescription of antibiotics for mild acute respiratory infections in children. *Bulletin of the Pan American Health Organisation* 1996;**30**(2):106–17.

#### Pelto 2004 *{published data only}*

Pelto GH, Santos I, Gonçalves H, Victora C, Martinez J, Habicht JP. Nutrition counseling training changes physician behavior and improves caregiver knowledge acquisition. *The Journal of Nutrition* 2004;**134**(2):357–62.

### Additional references

#### Baskett 2005

Baskett PJ, Nolan JP, Handley A, Soar J, Biarent D, Richmond S. European Resuscitation Council. European resuscitation council guidelines for resuscitation 2005. Section 9. Principles of training in resuscitation. *Resuscitation* 2005;**67**S1:S181–9.

#### Berkley 2005

Berkley JA, Maitland K, Mwangi I, Ngetsa C, Mwarumba S, Lowe BS, et al. Use of clinical syndromes to target antibiotic prescribing in seriously ill children in malaria endemic area: observational study. *BMJ* 2005;**330**(7498):995.

**Elie 2007**

Akl EA, Treweek S, Foy R, Francis J, Oxman AD, for the ReBEQI group. NorthStar, a support tool for the design and evaluation of quality improvement interventions in healthcare. *Implementation Science* 2007;**2**:19 doi:10.1186/1748-5908-2-19.

**EPOC 2007**

EPOC 2007. EPOC. Resources for review authors- reviewer tools. [www.epoc.cochrane.org/en/index.html](http://www.epoc.cochrane.org/en/index.html) (accessed 19 December 2007).

**Forsberg 2007**

Forsberg BC, Petzold MG, Thompson G, Allebeck P. Diarrhoea case management in low and middle-income countries-an unfinished agenda. *Bulletin of the World Health Organization* 2007;**85**(1):42-8.

**Forsetlund 2009**

Forsetlund L, Bjørndal A, Rashidian A, Jamtvedt G, O'Brien MA, Wolf F, et al. Continuing education meetings and workshops: effects on professional practice and health care outcomes. *Cochrane Database of Systematic Reviews* 2009, Issue 2. [DOI: 10.1002/14651858.CD003030.pub2]

**Gove 1999**

Gove S, Tamburlini G, Molyneux E, Whitesell P, Campbell H. Development and technical basis of simplified guidelines for emergency triage assessment and treatment in developing countries. WHO Integrated Management of Childhood Illness (IMCI) Referral Care Project. *Archives of Disease in Childhood* 1999;**81**(6):473-7.

**Higgins 2006**

Higgins JPT, Green S, (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 4.2.6 [Updated September 2006]. The Cochrane Collaboration, 2006. Available from [www.cochrane-handbook.org](http://www.cochrane-handbook.org).

**Irimu 2008**

Irimu G, Wamae A, Wasunna A, Were F, Ntoburi S, Opiyo N, et al. Developing and introducing evidence based clinical practice guidelines for serious illness in Kenya. *Archives of Disease in Childhood* 2008;**93**(9):799-804.

**Jabbour 1996**

Jabbour M, Osmond MH, Klassen TP. Life support courses: are they effective?. *Annals of Emergency Medicine* 1996;**28**:690-8.

**Jewkes 2003**

Jewkes F, Phillips B. Resuscitation training of paediatricians. *Archives of Disease in Childhood* 2003;**88**:118-21.

**McKenna 1985**

McKenna S, Glendon A. Occupational first aid training: decay in cardiopulmonary resuscitation (CPR) skills. *Journal of occupational psychology* 1985;**58**:109-17.

**Mello 2003**

Mello MJ, dos Santos ER, Buzzati R, et al. Pregnancy, childbirth, postpartum and newborn care guide for essential practice. In: Integrated management of childbirth. *Geneva: World Health Organization* 2003:1-178.

**Molyneux 2006**

Molyneux E, Ahmad S, Robertson A. Improved triage and emergency care for children reduces inpatient mortality in a resource-constrained setting. *Bulletin of the World Health Organization* 2006;**84**(4):314-9.

**Nolan 2001**

Nolan T, Angos P, Cunha AJ, Muhe L, Qazi S, Simoes EA, et al. Quality of hospital care for seriously ill children in less-developed countries. *Lancet* 2001;**357**(9250):106-10.

**Pio 2003**

Pio 2003. Standard case management of pneumonia in children in developing countries: the cornerstone of the acute respiratory infection programme. *Bulletin of the World Health Organization* 2003;**81**(4):298-300.

**Raupp 2007**

Raupp P, McCutcheon C. Neonatal resuscitation-An analysis of the transatlantic divide. *Resuscitation* 2007;**75**(2):345-9.

**Rowe 2002**

Rowe 2002. Design effects and intraclass correlation coefficients from a health facility cluster survey in Benin. *International Journal for Quality in Health Care* 2002;**14**(6):521-523.

**Rowe 2008**

Rowe AK, Rowe SY, Holloway KA, Ivanovska I, Muhe L, Lambrechts T. A systematic review of the effectiveness of shortening Integrated Management of Childhood Illness guidelines training: final report. World Health Organization 2008.

**Sazawal 2001**

Sazawal S, Black RE, for the Pneumonia Case Management Trials Group. Effect of pneumonia case management on mortality in neonates, infants, and preschool children: a meta-analysis of community-based trials. *Lancet Infectious Diseases* 2001;**3**(9):547-56.

**Sethi 2001**

Sethi D, Kwan I, Kelly AM, Roberts I, Bunn F. Advanced trauma life support training for ambulance crews. *Cochrane Database of Systematic Reviews* 2001, Issue 2. [DOI: 10.1002/14651858.CD003109]

**Shakiba 2003**

Shakiba H, Dinesh S, Anne MK. Advanced trauma life support training for hospital staff. *Cochrane Database of Systematic Reviews* 2003, Issue 3. [DOI: %3Chtml%3E%3Cbody id=%22body%22%3E10.1002/14651858.CD004173.pub2%3C/body%3E%3C/html%3E]

**Tulloch 1999**

Tulloch J. Integrated approach to child health in developing countries. *Lancet* 1999;**354** (Suppl. 2):S1116-20.

**WHO 2002**

WHO 2002. World Health Organization. Training course on the management of severe malnutrition. *Geneva: WHO* 2002.

**WHO 2005**

World Health Organization. The World Health Report: 2005: make every mother and child count. Geneva: World Health Organization 2005.

\* Indicates the major publication for the study



## CHARACTERISTICS OF STUDIES

### Characteristics of included studies [ordered by study ID]

#### Opiyo 2008

Methods	Randomised controlled trial (RCT)
Participants	Nurses Country: Kenya Phase 1: 83 nurses (28 intervention, 55 control) 97 practices in intervention group; 115 practices in control group Type of targeted behaviour - process of initiating newborn resuscitation
Interventions	Newborn resuscitation training (NRT) Duration of intervention - 1 day Co-intervention - self learning instruction manual provided to participants 2 weeks prior to training Control: standard practice Length of time during which intervention was measured after initiation of intervention - 50 days (phase 1)
Outcomes	Proportion of appropriate initial resuscitation steps Frequency of inappropriate/harmful practices Mortality
Notes	No difference between comparison groups in age and number of years worked Review authors also authors in the study (see conflict of interest statement) Primary analysis based on phase 1 data only Overall study quality: high risk of bias

#### *Risk of bias*

Item	Authors' judgement	Description
Adequate sequence generation?	Unclear	'...our intention was to randomise staff, stratified by place of work...'. The specific random approach was however not specified.
Allocation concealment?	Unclear	Health worker used as the unit of clustering but the random process incompletely reported.
Blinding? All outcomes	Yes	'The observers were blind to the training status of the health workers and were instructed not to try to ascertain health workers' training status...'
Contamination? All outcomes	Unclear	'We cannot exclude the possibility of cross-group contamination...'
Baseline characteristics?	Yes	'There were no significant differences in the ages...and years of experience between the groups...'

**Opiyo 2008** (Continued)

Incomplete data?	Yes	'32 allocated to intervention....28 providers observed', '58 allocated to control...55 providers observed'
Other risks of bias?	Yes	

**Senarath 2007**

Methods	Randomised controlled trial
Participants	Doctors, nurses, midwives Country: Sri Lanka 110 participants (59 intervention, 61 control) Type of targeted behaviour: general management - preparation and conduct of delivery care for newborn
Interventions	Essential newborn care (ENC) course Duration of intervention: 4 days Co-interventions: none Control: standard practice
Outcomes	Practices of essential newborn care at delivery
Notes	Reported results restricted to observation data only Length of time during which intervention was measured after initiation of intervention - 3 months 'Unit of analysis error present': unit of randomisation - hospitals; unit of analysis - observed delivery practices. Also, effects in experimental and control groups not directly compared. Overall study quality: high risk of bias

***Risk of bias***

Item	Authors' judgement	Description
Allocation concealment?	Yes	'...hospitals were randomly assigned to either of two groups...the intervention group...and control groups...'
Baseline outcome measurement? All outcomes	No	'There were some differences in the baseline level of practices between intervention and control groups...'
Blinding? All outcomes	No	'The principal investigator made observations in labor room...'
Contamination? All outcomes	Yes	Hospitals were randomly allocated to comparison groups, and it is unlikely that the control group received ENC training

Senarath 2007 (Continued)

Incomplete data?	Yes	'...participants (48 before and after the intervention) was selected...'. Effect of intervention on observed practices reported for 48 participants before and after the intervention'
Other risks of bias?	No	'Unit of analysis error'

### Characteristics of excluded studies *[ordered by study ID]*

Bryce 2005	Non-randomised design with concurrent health system strengthening activities (complex intervention)
El-Arifeen 2004	Data on referral rate for very ill children (outcome of interest) not reported
Gouws 2004	No baseline assessment of outcomes in IMCI trained and untrained groups
Nadel 2000	Study has a historical group only and used mock scenarios to assess practice
Ochoa 1996	Study did not include seriously ill children (only considered mild ARI episodes)
Pelto 2004	Study focused on an IMCI derived nutrition counselling protocol in apparently well children

## DATA AND ANALYSES

### Comparison 1. Opiyo 2008

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Mortality	1	90	Risk Ratio (M-H, Fixed, 95% CI)	0.77 [0.40, 1.48]

### Comparison 2. Senarath 2007


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Practice of preparedness of resuscitation	1	48	Mean Difference (IV, Fixed, 95% CI)	8.83 [6.55, 11.11]
2 Preparedness for resuscitation - adjusted for clustering	1		Mean Difference (Fixed, 95% CI)	8.83 [6.41, 11.25]

### Analysis 1.1. Comparison 1 Opiyo 2008, Outcome 1 Mortality.

Review: In-service training for health professionals to improve care of the seriously ill newborn or child in low and middle-income countries (Review)

Comparison: 1 Opiyo 2008

Outcome: 1 Mortality




## Analysis 2.1. Comparison 2 Senarath 2007, Outcome 1 Practice of preparedness of resuscitation.

Review: In-service training for health professionals to improve care of the seriously ill newborn or child in low and middle-income countries (Review)

Comparison: 2 Senarath 2007

Outcome: 1 Practice of preparedness of resuscitation




## Analysis 2.2. Comparison 2 Senarath 2007, Outcome 2 Preparedness for resuscitation - adjusted for clustering.

Review: In-service training for health professionals to improve care of the seriously ill newborn or child in low and middle-income countries (Review)

Comparison: 2 Senarath 2007

Outcome: 2 Preparedness for resuscitation - adjusted for clustering



## APPENDICES

### Appendix I. Detailed search strategy: CENTRAL / EPOC Register (up to May Week 1 2009)

#### Search terms

- #1 MeSH descriptor Inservice Training explode all trees
- #2 MeSH descriptor Health Personnel explode all trees with qualifier: ED
- #3 MeSH descriptor Internship and Residency, this term only
- #4 (staff or employee\* or clinician\* or physician? or nurs\* or midwife\* or midwiv\* or pharmacist\* or specialist\* or practitioner\* or dietician\* or nutritionist\*) NEXT (train\* or course\* or development or education or teach\*):ti or (staff or employee\* or clinician\* or physician? or nurs\* or midwife\* or midwiv\* or pharmacist\* or specialist\* or practitioner\* or dietician\* or nutritionist\*) NEXT (train\* or course\* or development or education or teach\*):ab
- #5 (inservice or in NEXT service or life NEXT support) NEAR/2 (train\* or course\* or development or education or teach\*):ti or (inservice or in NEXT service or life NEXT support) NEAR/2 (train\* or course\* or development or education or teach\*):ab
- #6 "on the job training":ti or "on the job training":ab
- #7 (#1 OR #2 OR #3 OR #4 OR #5 OR #6)
- #8 MeSH descriptor Case Management, this term only
- #9 MeSH descriptor Critical Care explode all trees
- #10 MeSH descriptor Life Support Care, this term only
- #11 MeSH descriptor Critical Illness, this term only
- #12 MeSH descriptor Acute Disease, this term only
- #13 MeSH descriptor Emergency Medical Services explode all trees
- #14 MeSH descriptor Emergency Medicine, this term only
- #15 MeSH descriptor Emergency Treatment explode all trees
- #16 MeSH descriptor Emergency Nursing, this term only
- #17 (case NEXT management):ti or (case NEXT management):ab
- #18 (emergency NEXT triage\*):ti or (emergency NEXT triage\*):ab
- #19 (life NEXT support):ti or (life NEXT support):ab
- #20 (resuscitation):ti or (resuscitation):ab
- #21 (first NEXT aid):ti or (first NEXT aid):ab
- #22 (referral or urgent) NEAR/2 care:ti or (referral or urgent) NEAR/2 care:ab
- #23 (critical\* or emergency or intensive or serious\* or sever\* or acute\*) NEAR/2 (care or ill or illness\* or treatment or therap\*):ti or (critical\* or emergency or intensive or serious\* or sever\* or acute\*) NEAR/2 (care or ill or illness\* or treatment or therap\*):ab
- #24 (#8 OR #9 OR #10 OR #11 OR #12 OR #13 OR #14 OR #15 OR #16 OR #17 OR #18 OR #19 OR #20 OR #21 OR #22 OR #23)
- #25 MeSH descriptor Child explode all trees
- #26 MeSH descriptor Infant explode all trees
- #27 MeSH descriptor Child Care explode all trees
- #28 MeSH descriptor Pediatrics explode all trees
- #29 MeSH descriptor Pediatric Nursing explode all trees
- #30 MeSH descriptor Perinatal Care, this term only
- #31 (child\* or infant\* or pediatric or paediatric or newborn\* or neonat\* or baby or babies or kid\* or toddler\*):ti or (child\* or infant\* or pediatric or paediatric or newborn\* or neonat\* or baby or babies or kid\* or toddler\*):ab
- #32 (#25 OR #26 OR #27 OR #28 OR #29 OR #30 OR #31)
- #33 MeSH descriptor Child Care explode all trees with qualifier: ED
- #34 MeSH descriptor Pediatrics explode all trees with qualifier: ED
- #35 MeSH descriptor Pediatric Nursing explode all trees with qualifier: ED
- #36 (#33 OR #34 OR #35)
- #37 MeSH descriptor Critical Care explode all trees with qualifier: ED
- #38 MeSH descriptor Life Support Care, this term only with qualifier: ED
- #39 MeSH descriptor Emergency Medical Services explode all trees with qualifier: ED
- #40 MeSH descriptor Emergency Medicine, this term only with qualifier: ED
- #41 MeSH descriptor Emergency Treatment explode all trees with qualifier: ED

- #42 MeSH descriptor Emergency Nursing, this term only with qualifier: ED  
 #43 (#37 OR #38 OR #39 OR #40 OR #41 OR #42)  
 #44 MeSH descriptor Intensive Care, Neonatal, this term only  
 #45 MeSH descriptor Diarrhea, Infantile, this term only  
 #46 MeSH descriptor Infant, Newborn, Diseases explode all trees  
 #47 "Acute Respiratory Infection" or "Acute Respiratory Infections":ti or "Acute Respiratory Infection" or "Acute Respiratory Infections":ab  
 #48 (#44 OR #45 OR #46 OR #47)  
 #49 "Control of Diarrheal Disease" or "Control of Diarrheal Diseases":ti or "Control of Diarrheal Disease" or "Control of Diarrheal Diseases":ab  
 #50 "Neonatal Resuscitation Program" or "Neonatal Resuscitation Programs":ti or "Neonatal Resuscitation Program" or "Neonatal Resuscitation Programs":ab  
 #51 "Essential Newborn Care":ti or "Essential Newborn Care":ab  
 #52 "Integrated Management of Childhood Illness":ti or "Integrated Management of Childhood Illness":ab  
 #53 (#49 OR #50 OR #51 OR #52)  
 #54 (#7 AND #24 AND #32)  
 #55 (#24 AND #36)  
 #56 (#32 AND #43)

## Appendix 2. Detailed search strategy: Ovid MEDLINE(R) 1950 to May Week 1 2009

### Search terms

1	exp Inservice Training/
2	exp Health Personnel/ed [Education]
3	"Internship and Residency"/
4	((staff or employee? or clinician? or physician? or nurs\$ or midwif\$ or midwiv\$ or pharmacist? or specialist? or practitioner? or dietician? or nutritionist?) adj (train\$ or course? or development or education or teach\$)).tw.
5	((inservice or in-service or life support) adj2 (train\$ or course? or development or education or teach\$)).tw.
6	on the job training.tw.
7	or/1-6
8	Case Management/
9	exp Critical Care/
10	Life Support Care/
11	Critical Illness/
12	Acute Disease/
13	exp Emergency Medical Services/

(Continued)

14	Emergency Medicine/
15	exp Emergency Treatment/
16	Emergency Nursing/
17	case management.tw.
18	emergency triage?.tw.
19	life support.tw.
20	resuscitation.tw.
21	first aid.tw.
22	((referral or urgent) adj2 care).tw.
23	((critical\$ or emergency or intensive or serious\$ or sever\$ or acute\$) adj2 (care or ill or illness\$ or treatment or therap\$)).tw.
24	or/8-23
25	exp Child/
26	exp Infant/
27	exp Child Care/
28	Pediatrics/
29	Neonatology/
30	Perinatology/
31	Pediatric Nursing/
32	Perinatal Care/
33	Neonatal Nursing/
34	(child\$ or infant? or pediatric or paediatric or newborn? or neonat\$ or baby or babies or kid? or toddler?).tw.
35	or/25-34
36	exp Child Care/ed [Education]
37	Pediatrics/ed [Education]



(Continued)

38	Neonatology/ed [Education]
39	Perinatology/ed [Education]
40	Pediatric Nursing/ed [Education]
41	Neonatal Nursing/ed [Education]
42	or/36-41
43	exp Critical Care/ed [Education]
44	Life Support Care/ed [Education]
45	exp Emergency Medical Services/ed [Education]
46	Emergency Medicine/ed [Education]
47	exp Emergency Treatment/ed [Education]
48	Emergency Nursing/ed [Education]
49	or/43-48
50	Intensive Care, Neonatal/
51	Diarrhea, Infantile/
52	Acute Respiratory Infection?.tw.
53	or/50-52
54	exp Infant, Newborn, Diseases/
55	Control of Diarrheal Disease?.tw.
56	Neonatal Resuscitation Program?.tw.
57	Essential Newborn Care.tw.
58	Integrated Management of Childhood Illness.tw.
59	or/55-58
60	7 and 24 and 35
61	24 and 42

(Continued)

62	35 and 49
63	7 and 53
64	7 and 24 and 54
65	or/59-64
66	Randomized Controlled Trial.pt.
67	random\$.tw.
68	intervention?.tw.
69	control\$.tw.
70	evaluat\$.tw.
71	effect\$.tw.
72	or/66-71
73	Animal/
74	Human/
75	73 not (73 and 74)
76	letter.pt.
77	editorial.pt.
78	comment.pt.
79	72 not (75 or 76 or 77 or 78)
80	65 and 79

### Appendix 3. Detailed search strategy: EMBASE (1980 to May Week 1 2009)

#### Search terms

1	In Service Training/
2	Staff Training/

(Continued)

3	Nurse Training/
4	Continuing Education/
5	Professional Development/
6	Medical Education/
7	Residency Education/
8	((staff or employee? or clinician? or physician? or nurs\$ or midwif\$ or midwiv\$ or pharmacist? or specialist? or practitioner? or dietician? or nutritionist?) adj (train\$ or course? or development or education or teach\$)).tw.
9	((inservice or in-service or life support) adj2 (train\$ or course? or development or education or teach\$)).tw.
10	on the job training.tw.
11	or/1-10
12	Case Management/
13	exp Intensive Care/
14	Critical Illness/
15	Disease Severity/
16	Acute Disease/
17	Injury Severity/
18	Emergency Medicine/
19	exp Emergency Treatment/
20	Emergency Nursing/
21	case management.tw.
22	emergency triage?.tw.
23	life support.tw.
24	resuscitation.tw.
25	first aid.tw.

(Continued)

26	((referral or urgent) adj2 care).tw.
27	((critical\$ or emergency or intensive or serious\$ or sever\$ or acute\$) adj2 (care or ill or illness\$ or treatment or therap\$)).tw.
28	or/12-27
29	exp Child/
30	exp Newborn/
31	exp Child Health Care/
32	exp Pediatrics/
33	exp Pediatric Nursing/
34	exp Postnatal Care/
35	Perinatal Care/
36	(child\$ or infant? or pediatric or paediatric or newborn? or neonat\$ or baby or babies or kid? or toddler?).tw.
37	or/29-36
38	Newborn Intensive Care/
39	Newborn Intensive Care Nursing/
40	Pediatric Intensive Care Nursing/
41	Pediatric Advanced Life Support/
42	Infantile Diarrhea/
43	Acute Respiratory Infection?.tw.
44	or/38-43
45	Emergency Medical Services Education/
46	exp Newborn Disease/
47	Control of Diarrheal Disease?.tw.
48	Neonatal Resuscitation Program?.tw.
49	Essential Newborn Care.tw.

(Continued)

50	Integrated Management of Childhood Illness.tw.
51	or/47-50
52	11 and 28 and 37
53	11 and 44
54	37 and 45
55	11 and 28 and 46
56	or/51-55
57	Randomized Controlled Trial/
58	Time Series Analysis/
59	random\$.tw.
60	experiment\$.tw.
61	(time adj series).tw.
62	(pre test or pretest or post test or posttest).tw.
63	impact.tw.
64	intervention?.tw.
65	chang\$.tw.
66	evaluat\$.tw.
67	effect?.tw.
68	compar\$.tw.
69	control\$.tw.
70	or/57-69
71	Nonhuman/
72	letter.pt.
73	editorial.pt.

(Continued)

74	70 not (71 or 72 or 73)
75	56 and 74

#### Appendix 4. Detailed search strategy: CINAHL (1982 to March Week 1 2008)

##### Search terms

1	exp Staff Development/
2	exp Health Personnel/ed [Education]
3	"Internship and Residency"/
4	((staff or employee? or clinician? or physician? or nurs\$ or midwif\$ or midwiv\$ or pharmacist? or specialist? or practitioner? or dietician? or nutritionist?) adj (train\$ or course? or development or education or teach\$)).tw.
5	((inservice or in-service or life support) adj2 (train\$ or course? or development or education or teach\$)).tw.
6	on the job training.tw.
7	or/1-6
8	Case Management/
9	exp Critical Care/
10	Life Support Care/
11	Critical Illness/
12	Acute Disease/
13	exp Emergency Medical Services/
14	Emergency Medicine/
15	First Aid/
16	exp Resuscitation/
17	exp Emergency Nursing/

(Continued)

18	case management.tw.
19	emergency triage?.tw.
20	life support.tw.
21	resuscitation.tw.
22	first aid.tw.
23	((referral or urgent) adj2 care).tw.
24	((critical\$ or emergency or intensive or serious\$ or sever\$ or acute\$) adj2 (care or ill or illness\$ or treatment or therap\$)).tw.
25	or/8-24
26	exp Child/
27	exp Child Care/
28	Child Health/
29	(child\$ or infant? or pediatric or paediatric or newborn? or neonat\$ or baby or babies or kid? or toddler?).tw.
30	Pediatrics/
31	Neonatology/
32	Perinatal Care/
33	Prenatal Care/
34	exp Pediatric Care/
35	exp Pediatric Nursing/
36	or/26-35
37	exp Child Care/ed [Education]
38	Child Health/ed [Education]
39	exp Pediatrics/ed [Education]
40	Perinatal Care/ed [Education]
41	Prenatal Care/ed [Education]

(Continued)

42	exp Pediatric Care/ed [Education]
43	exp Pediatric Nursing/ed [Education]
44	or/37-43
45	exp Critical Care/ed [Education]
46	Life Support Care/ed [Education]
47	exp Emergency Medical Services/ed [Education]
48	Emergency Medicine/ed [Education]
49	First Aid/ed [Education]
50	exp Resuscitation/ed [Education]
51	exp Emergency Nursing/ed [Education]
52	or/45-51
53	exp Intensive Care, Neonatal/
54	Acute Respiratory Infection?.tw.
55	or/53-54
56	exp Infant, Newborn, Diseases/
57	Control of Diarrheal Disease?.tw.
58	Neonatal Resuscitation Program?.tw.
59	Integrated Management of Childhood Illness.tw.
60	Pediatric Advanced Life Support/ed [Education]
61	or/57-60
62	7 and 25 and 36
63	25 and 44
64	36 and 52
65	7 and 55



(Continued)

66	7 and 25 and 56
67	or/61-66
68	Clinical Trial/
69	exp Pretest-Posttest Design/
70	exp Quasi-Experimental Studies/
71	Comparative Studies/
72	control\$.tw.
73	random\$.tw.
74	experiment\$.tw.
75	(time adj series).tw.
76	impact.tw.
77	intervention?.tw.
78	evaluat\$.tw.
79	effect\$.tw.
80	(pre test or pretest or post test or posttest).tw.
81	or/68-80
82	67 and 81

## Appendix 5. Detailed search strategy: LILACS (up to October 2008)

### Search terms

(inservice and training) or (inservice and course\$) or (inservice and workshop\$) or (inservice and education) or (inservice and program\$) or (capacitación and servicio) or (capacitação and serviço) [Palavras]

And

child or children or niño or criança or infant or infants or lactante or lactente or pediatric\$ or paediatric\$ or pediatria or pediatria or newborn or (recién and nacidos) or (recém and nascidos) or neonat\$ or baby or babies or kid or kids or toddler\$ [Palavras]

## Appendix 6. Detailed search strategy: ERIC (up to October 2008)

### Search terms

(DE=Inservice Education or DE=On the Job Training or TI=inservice training or TI=on the job training or TI=inservice course\* or TI=inservice workshop\* or TI=inservice education or TI=inservice program\* or TI=in service training or TI=in service course\* or TI=in service workshop\* or TI=in service education or TI=in service program\* or AB=inservice training or AB=on the job training or AB=inservice course\* or AB=inservice workshop\* or AB=inservice education or AB=inservice program\* or AB=in service training or AB=in service course\* or AB=in service workshop\* or AB=in service education or AB=in service program\*)

AND

(DE=Crisis Management or DE=Crisis Intervention or DE=Emergency Programs or DE=First Aid or TI=crisis management or TI=crisis intervention\* or TI=emergency program\* or AB=crisis management or AB=crisis intervention\* or AB=emergency program\* or TI=critical care or TI=critical\* ill\* or TI=critical treatment\* or TI=critical therap\* or TI=emergency care or TI=emergency ill\* or TI=emergency treatment\* or TI=emergency therap\* or TI=intensive care or TI=intensive ill\* or TI=intensive treatment\* or TI=intensive therap\* or TI=serious care or TI=serious ill\* or TI=serious treatment\* or TI=serious therap\* or TI=sever\* care or TI=sever\* ill\* or TI=sever\* treatment\* or TI=sever\* therap\* or TI=acute\* care or TI=acute\* ill\* or TI=acute\* treatment\* or TI=acute\* therap\* or TI=first aid or TI=life support or TI=urgent care or TI=resuscitation or AB=critical care or AB=critical\* ill\* or AB=critical treatment\* or AB=critical therap\* or AB=emergency care or AB=emergency ill\* or AB=emergency treatment\* or AB=emergency therap\* or AB=intensive care or AB=intensive ill\* or AB=intensive treatment\* or AB=intensive therap\* or AB=serious care or AB=serious ill\* or AB=serious treatment\* or AB=serious therap\* or AB=sever\* care or AB=sever\* ill\* or AB=sever\* treatment\* or AB=sever\* therap\* or AB=acute\* care or AB=acute\* ill\* or AB=acute\* treatment\* or AB=acute\* therap\* or AB=first aid or AB=life support or AB=urgent care or AB=resuscitation)

AND

(DE=Hospitalized Children or DE=children or DE=young children or DE=toddlers or DE=infants or DE=infant care or DE=premature infants or DE=neonates or DE=pediatrics or TI=child or TI=children or TI=infant or TI=infants or TI=pediatric\* or TI=paediatric\* or TI=newborn or TI=neonat\* or TI=baby or TI=babies or TI=kid or TI=kids or TI=toddler\* or AB=child or AB=children or AB=infant or AB=infants or AB=pediatric\* or AB=paediatric\* or AB=newborn or AB=neonat\* or AB=baby or AB=babies or AB=kid or AB=kids or AB=toddler\*)

## Appendix 7. Detailed search strategy: WHOLIS (up to October 2008)

### Search terms

words or phrase

inservice or 'in service' or 'on the job'

AND words or phrase

training or course\$ or education or workshop\$ or program\$

AND words or phrase

child\$ or infant\$ or pediatric\$ or paediatric\$ or newborn\$ or neonat\$ or baby or babies or kid or kids or toddler\$

## WHAT'S NEW

Last assessed as up-to-date: 21 May 2009.

22 March 2010	Amended	Minor edits
---------------	---------	-------------

## HISTORY

Protocol first published: Issue 2, 2008

Review first published: Issue 4, 2010

## CONTRIBUTIONS OF AUTHORS

NO and ME wrote the protocol, screened records for eligibility, extracted data, assessed methodological quality of included studies, interpreted findings and wrote the review. NO prepared the first draft of the review. ME commented on the manuscript.

## DECLARATIONS OF INTEREST

NO and ME are authors of one of the studies [Opiyo 2008](#) included in this review. The methodological quality of this study was also considered by another EPOC reviewer.

## SOURCES OF SUPPORT

### Internal sources

- KEMRI/Wellcome Trust Research Programme, Kenya.

### External sources

- South African Cochrane Centre, South Africa.
- Cochrane Effective Practice and Organisation of Care Group (EPOC), Norway.

## INDEX TERMS

### Medical Subject Headings (MeSH)

\*Developing Countries; Infant, Newborn; Inservice Training [\*methods]; Neonatology [\*education]; Perinatology [\*education]; Quality of Health Care [\*standards]; Randomized Controlled Trials as Topic

### MeSH check words

Child; Humans